

*Do you suffer  
from migraine headaches?*

*Are you currently taking  
immediate-release topiramate?*

Some important questions to  
ask yourself about your current  
migraine prevention treatment plan:

1. Are you currently taking immediate-release  
topiramate?

Yes  No

2. Do you sometimes miss one of your  
doses of medicine?

Yes  No

3. Are you experiencing side effects?

Yes  No

4. Are you satisfied with your current migraine  
prevention treatment?

Yes  No

FILL OUT  
& DISCUSS  
WITH YOUR  
DOCTOR!

See page 2 for topiramate  
SAVINGS AND SUPPORT PROGRAM

\$0\*

CO-PAY

# Patient savings and support with Trokendi XR® \$0 Co-pay Program\*

\$0  
CO-PAY

**Trokendi XR® Patient Savings and Support Program**

**PAY AS LITTLE AS \$0\***  
per prescription for your next 12 prescriptions\*

Call 1-866-398-0833 to activate your card or visit [www.TrokendiXR.com](http://www.TrokendiXR.com) to enroll.

**Trokendi XR**  
topiramate extended-release capsules  
200 mg 100 mg 50 mg

RxBIN: 610524  
RxCPCN: Loyalty  
ISSUER: (80840)  
RxGRP: 50777445  
ID: XXXXXXXX

\*Restrictions apply. See back offer for details.

DOWNLOAD CO-PAY CARD

## INDICATION

Trokendi XR (topiramate) extended-release capsules are used to prevent migraine headaches in adults and adolescents 12 years of age and older.

## Do not take Trokendi XR if

- You have recently consumed or plan to consume alcohol (i.e., within 6 hours prior to and 6 hours after Trokendi XR use)

## How do I get started?

**To the Patient:** You must present this card to the pharmacist along with your prescription to participate in this program. If you have any questions regarding your eligibility or benefits, or if you wish to discontinue your participation, call the Trokendi XR® program at 1-866-398-0833 (8:00AM-6:00PM EST, Monday-Friday). When you use this card, you are certifying that you understand the program rules, regulations, and terms and conditions. You are not eligible if prescriptions are paid by any state or other federally funded programs, including, but not limited to Medicare or Medicaid, Medicaid, VA or OIG or Tricare, or where prohibited by law; and you will otherwise comply with the terms above.

**To the Pharmacist:** When you use this card, you are certifying that you have not submitted and will not submit a claim for reimbursement under any federal, state or other governmental programs for this prescription.

- Submit transaction to McKesson Corporation using BIN #65024
- If primary commercial prescription insurance exists, input card information as secondary coverage and transmit using the COB segment of the NCPDP transaction. Applicable discounts will be displayed in the transaction response.
- Acceptance of this card and your submission of claims for the Trokendi XR® program are subject to the LoyaltyCard® program Terms and Conditions posted at [www.mckesson.com/loyalty](http://www.mckesson.com/loyalty)
- Patient is not eligible if prescriptions are paid in part or full by any state or federally funded programs, including but not limited to Medicare or Medicaid, Medicaid, VA, OIG or Tricare and where prohibited by law.
- For questions regarding setup, claim transmission, patient eligibility or other issues, call the LoyaltyCard® for Trokendi XR® program at 877-264-2448 (8:00 AM-8:00 PM EST, Monday-Friday).

Powered By  
**MCKESSON**

Supernus reserves the right to rescind, revoke, or amend this offer without notice at any time.

**Supernus** Trokendi XR is a registered trademark of Supernus Pharmaceuticals, Inc.  
© 2015 Supernus Pharmaceuticals, Inc. All rights reserved. SPN.TRO.2015-0225

**Trokendi XR**  
topiramate extended-release capsules  
200 mg 100 mg 50 mg

\*For full terms and conditions, please see the Trokendi XR Co-pay Card, or visit [www.TrokendiXR.com](http://www.TrokendiXR.com).

ONCE-A-DAY FOR MIGRAINE PREVENTION

**Trokendi XR**  
(topiramate) extended-release capsules



Capsules shown are not actual size.

Please refer to the Important Safety Information on page 3, or the Medication Guide available at [www.TrokendiXR.com](http://www.TrokendiXR.com) for more information on Trokendi XR.

## Trokendi XR® (topiramate) extended-release capsules, for oral use

### INDICATION

Trokendi XR® (topiramate) extended-release capsules are used to prevent migraine headaches in adults and adolescents 12 years and older.

### IMPORTANT SAFETY INFORMATION

#### Who should not take Trokendi XR?

Do not take Trokendi XR if you have recently consumed or plan to consume alcohol (i.e., within 6 hours prior to and 6 hours after Trokendi XR use)

**Swallow Trokendi XR capsules whole. Do not sprinkle on food, chew, or crush.**

#### What are the possible side effects of Trokendi XR?

##### Trokendi XR can cause serious side effects, including: Eye problems.

Serious eye problems include sudden decrease in vision with or without eye pain or redness, a blockage of fluid that may cause increased pressure in the eye (secondary angle closure glaucoma). Call your healthcare provider right away if you have new eye symptoms, including any new problems with your vision.

**Decreased sweating and increased body temperature (fever).** People, especially children, should be watched for signs of decreased sweating and fever, especially in hot temperatures. Some people may need to be hospitalized for this condition.

**Increased levels of acid in the blood (metabolic acidosis).** If left untreated, metabolic acidosis can cause brittle or soft bones (osteoporosis, osteomalacia, osteopenia), kidney stones, can slow the rate of growth in children, and may possibly harm the unborn child of pregnant patients.

**High levels of ammonia in the blood.** High ammonia in the blood can affect mental activities, slow alertness, cause tiredness, or cause vomiting. Blood ammonia levels have been shown to rise when Trokendi XR is taken with a medicine called valproic acid (e.g., DEPAKENE® and DEPAKOTE®).

**Kidney stones.** Drink plenty of fluids when taking Trokendi XR to decrease your chances of getting kidney stones.

**Low body temperature.** Taking Trokendi XR when you are also taking valproic acid may cause a drop in body temperature to less than 95°F, tiredness, confusion, or coma.

**Effects on thinking and alertness.** Trokendi XR may affect how you think, and can cause confusion and problems with concentration, attention, memory, or speech. Trokendi XR may cause depression or mood problems, tiredness, and sleepiness.

**Dizziness or loss of muscle coordination.**

**The most common side effects include** tingling of the arms and legs (paresthesia), not feeling hungry, nausea, weight loss, abnormal vision, a change in the way foods taste, nervousness, speech problems, dizziness, slow reactions, upper respiratory tract infection, sleepiness, diarrhea, pain in abdomen and difficulty with memory. These are not all the possible side effects of Trokendi XR. For more information, ask your healthcare provider or pharmacist.

**Like other antiepileptic drugs, Trokendi XR may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.** Before you take Trokendi XR, tell your healthcare provider if you have or have had depression, mood problems, or suicidal thoughts or behavior. Call a

healthcare provider right away if you have thoughts about suicide or dying; have attempted to commit suicide; have new or worsening depression or anxiety; feel agitated or restless; experience panic attacks, trouble sleeping (insomnia), or new or worsening irritability; feel or act more aggressive, angry, or violent; act on dangerous impulses; have an extreme increase in activity and talking (mania); or experience other unusual changes in behavior or mood.

**Before taking Trokendi XR, tell your healthcare provider about any other medical conditions,** including if you have kidney problems, kidney stones, or are getting kidney dialysis; have a history of metabolic acidosis (too much acid in the blood); have liver problems; have weak, brittle or soft bones (osteomalacia, osteoporosis, osteopenia, or decreased bone density); have lung or breathing problems; have eye problems, especially glaucoma; have diarrhea; have a growth problem; are on a diet high in fat and low in carbohydrates, which is called a ketogenic diet; are having surgery; are pregnant or plan to become pregnant; or if you are breastfeeding. Trokendi XR passes into your breast milk. Breastfed babies may be sleepy or have diarrhea. Talk to your healthcare provider about the best way to feed your baby if you take Trokendi XR.

**Trokendi XR can harm your unborn baby.** If you take Trokendi XR during pregnancy, your baby has a higher risk for the birth defects of cleft lip, cleft palate, and being smaller than expected at birth. These defects can begin early in pregnancy, even before you know you are pregnant. The long term effects of this are unknown.

**Tell your healthcare provider about any other medicines you take,** including prescription and nonprescription medicines, vitamins, and herbal supplements. Trokendi XR and other medicines may affect each other, causing side effects. Especially tell your healthcare provider if you take valproic acid (e.g., DEPAKENE or DEPAKOTE); any medicines that impair or decrease your thinking, concentration, or muscle coordination; or birth control pills. Trokendi XR may make your birth control pills less effective.

**Do not stop Trokendi XR without first talking to a healthcare provider.** If you have epilepsy and you stop taking Trokendi XR suddenly, you may have seizures that do not stop. Your healthcare provider will tell you how to stop taking Trokendi XR slowly.

**Do not drive a car or operate heavy machinery until you know how Trokendi XR affects you.** Trokendi XR can slow your thinking and motor skills, and may affect vision.

**Please refer to the Medication Guide for additional important patient information.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

ONCE-A-DAY FOR MIGRAINE PREVENTION

**Trokendi XR**®  
(topiramate) extended-release capsules



25 mg

50 mg

100 mg

200 mg

Capsules shown are not actual size.